

原著

Hiroharu Mifune, Yoshihiro Nishi, Yuji Tajiri, Akira Yabuki

A-type natriuretic peptide level in angiotensin II type 1a receptor knockout mice.
Experimental Animals 61(2) : 131-8, 2012.

Hiroh Kaku, Yuji Tajiri, Kentaro Yamada

Anorexigenic Effects of Miglitol in Concert with the Alterations of Gut Hormone
Secretion and Gastric Emptying in Healthy Subjects
Horm Metab Res 44(4) : 312-8, 2012.

Masanori Ohta, Yuji Tajiri, Hiroshi Yamato, Masaharu Ikeda

Effects of exercise therapy alone and in combination with a calcium channel blocker
or an angiotensin receptor blocker in hypertensive patients.
Clinical and Experimental Hypertension 34(7) : 523-9, 2012.

Naokazu Hamada, Yoshihiro Nishi, Yuji Tajiri, Kentaro Setoyama, Ryoza Kamimura,
Kenkichi Miyahara, Norihito Nuruki, Hiroshi Hosoda, Kenji Kanagawa,
Masayasu Kojima, Hiroharu Mifune

Disrupted Regulation of Ghrelin Production Under Antihypertensive Treatment in
Spontaneously Hypertensive Rats
Circulation Journal 76(6) : 1423-9, 2012.

Hiroharu Mifune, Yoshihiro Nishi, Yuji Tajiri, Taku Masuyama, Hiroshi Hosoda,
Kenji Kanagawa, Masayasu Kojima

Increased production of active ghrelin is relevant to hyperphagia in nonobese
spontaneously diabetic Torii rats
Metabolism 61(4) : 491-5, 2012.

Shuichi Otabe, Nobuhiko Wada, Toshihiko Hashinaga, Xiaohoug Yuan, Isao Shimokawa,
Tomoka Fukutani, Kayo Tanaka, Tsuyoshi Ohki, Satomi Kakino, Yayoi Kurita,
Hitomi Nakayama, Yuji Tajiri and Kentaro Yamada

Hyperadiponectinemia protects against premature death in metabolic syndrome
model mice by inhibiting AKT signaling and chronic inflammation
Journal of Endocrinology 213(1) : 67-76, 2012.

Hiroharu Mifune, Yoshihiro Nishi, Yuji Tajiri and Akira Yabuki

Different A-Type Natriuretic Peptide Level in Five Strains of Mice

The Journal of Veterinary Medical Science 74(4) : 499-502,2012.

Ohta M, Tajiri Y, Yamamoto H, Ikeda M

Effects of exercise therapy alone and in combination with a calcium channel blocker or an Angiotensin receptor blocker in hypertensive patients

Clinical and Experimental Hypertension 34(7) : 523-9,2012.

Tsuyoshi Ohki, Yoshifumi Sato, Tatsuya Yoshizawa, Ken-ichi Yamamura,

Kentaro Yamada, Kazuya Yamagata

Identification of hepatocyte growth factor activator (Hgfac) gene as a target of HNF1 α in mouse β -cells

Biochemical and Biophysical Research Communications 425(3) : 619-24,2012.

Hayato Kaida, Seiji Kurata, Akihiko Kawahara, Yuji Hiromatsu, Masayoshi Kage,

Masatoshi Ishibashi

High ¹⁸F-FDG uptake in sporadic paraganglioma of the retroperitoneum may be related to intra-tumor haemorrhage and macrophages

Hellenic Journal of Nuclear Medicine 15(3) : 261,2012.

Shuichi Yatsuga, Yuji Hiromatsu, Shigekazu Sasaki, Hirotohi Nakamura,

Koju Katayama, Junko Nishioka and Yasutoshi Koga

A two-day-old hyperthyroid neonate with thyroid hormone resistance born to a mother with well-controlled Graves' disease : a case report

Journal of Medical Case Reports 6(1) : 246,2012.

Yoriko Hara, Sanae Iwashita, Kouzabu Ishii, Chizuko Inada, Akira Okada, Yuji Tajiri,

Hitomi Nakayama, Tomoko Kato, Keiko Nishida, Yasuhiro Ogata, Hisamitsu Omori,

Toru Morinaga, Midori Yamaguchi, Motoyuki Nakao, Koji Tsuboi, Raoul Breugelmans,

Yoko Ishihara

The reliability and validity of the Japanese version of the Diabetes Family Behavior Checklist (DFBC) for assessing the relationship between Type 2 diabetes mellitus patients and their families with respect to adherence to treatment regimen

Diabetes Research and Clinical Practice 99(1) : 39-47,2013.

Yuji Tajiri, Koji Yonemoto, Kentaro Yamada

Subcutaneous fat of extremities independent of abdominal fat does not promote insulin resistance and cardiovascular risk in Japanese type 2 diabetic patients.

Diabetology International 4(1) : 45-51,2013.

江口洋幸, 南泰山, 佐藤秀一, 谷淳一, 廣松雄治

頸部超音波検査を契機に発見された Zenker 憩室の 1 例

内科 110(2) : 325-7, 2012.